

Discontinue Use of Central Line?	✓ Central Line Care Protocol implemented?
Discontinue Use of Indwelling Urinary Catheter?	✓ Indwelling Urinary Catheter Care Protocol implemented?
Discontinue Use of Ventilator?	✓ VAE Prevention Protocols implemented?
Patient at Risk for Agitation or Delirium?	✓ Pain Management: Optimal use of pharmacologic and non-pharmacologic pain management interventions?
	✓ Agitation Management: Optimal use of pharmacologic and non-pharmacologic agitation prevention and management interventions?
	✓ Delirium Management: Optimal use of pharmacologic and non-pharmacologic delirium prevention and management interventions?
Patient at Risk for Falls?	✓ Falls Prevention Protocol implemented?
Patient at Risk for Pressure Ulcers?	✓ Pressure Ulcer Protocol implemented?
Patient at Risk for Venous Thromboembolism (VTE)?	✓ VTE Prevention Protocol implemented?
Patient at Risk for a Change in Condition?	✓ Are there any new or unexpected signs or symptoms related to clinical deterioration (e.g., sepsis, <i>Clostridium difficile</i> , delirium, surgical site infection, any decompensation or reason for rapid response activation)?
Medication Reconciliation Completed?	✓ Was medication reconciliation completed upon admission to and upon discharge from the critical care unit?
Readmission Prevention Provisions Completed?	✓ Referrals to social work or case management done?
	✓ Assessment conducted and action taken for palliative care or hospice?
Patient-Centered Care Considerations:	✓ Have opportunities been actively provided to include patients and caregivers as participatory members of the care team?
	✓ Are the patient and family being included in developing the goal(s) of care and post-hospital care needs?