

Discontinue Use of Central Line?	✓ Central Line Care Protocol implemented?
Discontinue Use of Indwelling Urinary Catheter?	✓ Indwelling Urinary Catheter Care Protocol implemented?
Patient at Risk for Falls?	✓ Falls Prevention Protocol implemented?
Patient at Risk for Pressure Ulcers?	✓ Pressure Ulcer Protocol implemented?
Patient at Risk for VTE?	✓ VTE Prevention Protocol implemented?
Patient at Risk for Decompensation or Complication?	✓ Are there any new or unexpected signs or symptoms related to the patient's diagnosis or other illnesses (e.g., sepsis, <i>Clostridium difficile</i> , delirium, surgical site infection, or any decompensation or reason for rapid response activation)?
Patient at Risk for Readmission?	✓ Were appropriate referrals made to prevent readmissions, such as physical therapy, social work, nutrition?
	✓ Were patient and family concerns addressed through ongoing education and teach-back from multi-disciplinary staff in preparation for self-care at discharge?
	✓ Was a pharmacist consulted if patient is taking any high-risk medications (insulin, anticoagulants, and opioids)?
Patient-Centered Care Considerations:	✓ Have opportunities been actively provided to include patients and caregivers as participatory members of the care team?
	✓ Are the patient and family being included in developing the goal(s) of care and post-hospital care needs?
Day Before or Day of Anticipated Discharge:	✓ Discharge medication reconciliation completed?
	✓ All medication reconciliation discrepancies resolved?
	✓ Education provided on high-risk medications (insulin, anticoagulants, and opioids)?
	✓ Has there been coordination and communication with post-hospital care providers?
	✓ Do patients and families understand discharge instructions?